LIVER LOVE

Clinical Applications

- Provides Micronutrients, Phytonutrients, and Cofactors that Support Detoxification of Xenobiotics and Xenoestrogens*
- Supports Healthy Estrogen Metabolism*
- Supports Antioxidant Mechanism and Glutathione Production*

Liver Love is a comprehensive formula designed to support phase I and phase II liver detoxification of environmental pollutants, endocrine disruptors, estrogen metabolites, xenoestrogens, and other toxins. Liver Love also supports antioxidant activity throughout the detoxification process. Micronutrients, phytonutrients, and activated cofactors provide additional support for energy production, cellular protection, and liver function during crucial metabolic biotransformation processes.*

All MSW Nutrition Formulas Meet or Exceed cGMP Quality Standards

Discussion

Xenobiotics (chemicals foreign to a living organism) have the potential to disrupt metabolism and negatively affect cellular health.^[1-3] Classes of xenobiotics include pesticides, petroleum-based plastic compounds, industrial chemicals, and xenoestrogens. Liver Love comprises an array of compounds to support detoxification and elimination of these potentially toxic molecules. Man-made xenoestrogens (including BPA, DDT, and DES), act as endocrine disruptors and can alter hormonal function in sensitive tissues including breast, uterus, cervix, and prostate.^[4,5] Xenoestrogens at very low levels are believed to disrupt neurotransmitter balance, glucose homeostasis, normal reproduction, and healthy metabolism.^[5] Detoxification of xenobiotics is a complex process that requires micronutrients, phytonutrients, energy, and adequate antioxidant support for safe and effective completion.^{*[6]}

Antioxidant and Detoxification Support Several nutrients support antioxidant activity, both phases of detoxification, and the health and function of the liver (the major site of detoxification). Milk thistle extract contains silymarin, a compound found to limit the entry of hepatotoxins, donate sulfhydryl groups for detoxification, and increase hepatic glutathione by over 35%.^[7] Its action in the liver reduces fat peroxidation and fibrous tissue formation, supports a normal immune and inflammatory response, promotes protein synthesis and tissue regeneration, and supports glucuronidation and glutathione levels.^[8] Alpha-lipoic acid is both water- and fat-soluble. It supports glutathione metabolism, helps regenerate antioxidant vitamins C and E, and helps maintain the ratio of reduced-to-oxidized CoQ10 in mitochondria.^[7] The redox couple of lipoic acid and dihydrolipoic acid stabilizes NF-kappaB transcription and may help support healthy immune functions in the body.^[9,10] Methylselenocysteine (MSC) is considered a well-tolerated form of the trace element selenium and may support normal cell-life regulation.^[11] Selenium provides antioxidant support via glutathione in the body, which, in turn, is incorporated into crucial antioxidant and detoxification enzymes. Glutathione supports antioxidant activity, phase II detoxification, and the normal breakdown of metabolites, toxins, and other compounds. NAC supports phase II sulfation reactions as well.^[7] Calcium D-glucarate has been added to support glucuronidation. 5-methyltetrahydrofolate (5-MTHF) is present as Quatrefolic[®] (a stable, bioavailable form of folate) to support methylation, energy generation, and phase I and phase II activity.*

Phytonutrients A variety of phytonutrients support antioxidant activity in the body. **Green tea catechins** have been found to assist in free-radical scavenging, support detoxification through modification of phase I and phase II enzymes, and support normal cell-life regulation via multiple signaling pathways.^[13,14] Bioflavonoids, including **resveratrol, quercetin**, and the highly absorbable FlavitPURE[™] form of **dihydroquercetin (DHQ)**, support phase I detoxification as well as intermediary antioxidant protection.^[1] **Pterostilbene**, a highly absorbable, methylated form of resveratrol, is thought to work together with quercetin in supporting normal cell-life regulation.^[15] Turmeric extract provides curcumin, a phytonutrient valued for its promotion of antioxidant activity, support of metabolic detoxification, and modulation of cytokine production.^[16,17] BioPerine[®], a patented form of piperine from **black pepper**, has been added to enhance the absorption of nutrients, particularly curcumin.^{*[18]}

Xenoestrogen Metabolism Liver Love provides diindolymethane (DIM) and glucoraphanin as SGS[™]. DIM promotes healthy estrogen metabolism and creates a better balance of estrogen metabolites (2-OH, 4-OH, 16-alpha-OH) through phase I cytochrome P450 enzyme induction and promotion of 2-hydroxylation.^[19,20] The action of DIM is complemented by glucoraphanin, which supports long-term antioxidant activity and phase II detoxification of less-desirable estrogen metabolites and xenoestrogens.^[21,22] Glucoraphanin and its metabolite sulforaphane are found to be effective, long-acting, indirect antioxidants and significant inducers of phase II detoxification enzymes.^[23,24] These actions may help support healthy estrogen balance and may be crucial for the health of estrogen-sensitive tissue.*



Supplement Facts

Serving Size: 2 Capsules Servings Per Container: 60

	Amount Per Serving	%DV
Folate (as Quatrefolic [®] (6S)-5-methyltetrahydrofolic acid, glucosamine salt)	200 mcg DFE	50%
Calcium (as tricalcium phosphate and calcium d-glucarate tetrahydrate)	75 mg	6%
Selenium (as methylselenocysteine)	15 mcg	27%
Calcium D-Glucarate Tetrahydrate	250 mg	**
Green Tea Aqueous Extract (Camellia sinensis)(leaf)	250 mg	**
(80% polyphenols, 60% catechins, 30% EGCG, 6% caffeine)		
Alpha-Lipoic Acid	100 mg	**
N-Acetyl-L-Cysteine	100 mg	**
Milk Thistle Extract (Silybum marianum)(seed)(30% silybins)	100 mg	**
DIM (diindolylmethane)	75 mg	**
Quercetin (as quercetin dihydrate)(from Sophora japonica)(bud)	50 mg	**
Turmeric Extract (Curcuma longa)(rhizome)(95% curcuminoids)	50 mg	**
trans-Resveratrol (as Polygonum cuspidatum root extract)	18.5 mg	**
pTeroPure® <i>trans</i> -Pterostilbene	15.5 mg	**
truebroc® Glucoraphanin (from broccoli extract)(Brassica oleracea italica)(see	l) 15 mg	**
Dihydroquercetin	5 mg	**
BioPerine® Black Pepper Extract (Piper nigrum)(fruit)	5 mg	**
** Daily Value (DV) not established.		

Other Ingredients: HPMC (capsule), stearic acid, magnesium stearate, and silica.

Quatrefolic[®] is a registered trademark of Gnosis S.p.A. Produced under US Patent 7,947,662.

Produced under US patent 6,521,818 licensed from Brassica Protection Products LLC; truebroc is a registered trademark of Brassica Protection Products LLC. true

pTeroPure is a trademark of ChromaDex, Inc.

BioPerine is a registered trademark of Sabinsa Corp. BioPerine is protected by US patents 5,536,506; 5,744,161; 5.972.382; and 6.054.585

References

1. Gaby AR. Nutritional Medicine. Concord, NH: Fritz Perlberg Publishing; 2011.

2. Colborn T, vom Saal FS, Soto AM. Developmental effects of endocrine-disruptingchemicals in wildlife and humans. Environ Health Perspect. 1993 Oct;101(5):378-84. Review. [PMID: 80805061

3. Alexander BJ, Ames BN, Baker SM, et al. Textbook of Functional Medicine. Gig Harbor, WA: The Institute for Functional Medicine; 2010.

4. Singleton DW, Khan SA. Xenoestrogen exposure and mechanisms of endocrine disruption. Front Biosci. 2003 Jan 1;8:s110-8. Review. [PMID: 12456297] 5. Nadal A, Ropero AB, Laribi O, et al. Nongenomic actions of estrogens and xenoestrogens by binding at a plasma membrane receptor unrelated to estrogen receptor alpha and estrogen receptor beta. *Proc Natl Acad* Sci U S A. 2000 Oct 10:97(21):11603-8. [PMID: 11027358]

6. Liska DJ. The detoxification enzyme systems. Altern Med Rev. 1998 Jun;3(3):187-98. Review. [PMID: 9630736]

7. Krinsky DL, LaValle JB, Hawkins EB, et al. Natural Therapeutics Pocket Guide. 2nd ed. Hudson, OH: Lexi-Comp; 2003. 8. Pradhan SC, Girish C. Hepatoprotective herbal drug, silymarin from experimental pharmacology to clinical medicine. Indian J Med Res. 2006 Nov;124(5):491-504. [PMID: 17213517] 9. Suzuki YJ, Aggarwal BB, Packer L. Alpha-lipoic acid is a potent inhibitor of NFkappa B activation in human T cells. Biochem Biophys Res Commun. 1992 Dec 30;189(3):1709-15. [PMID: 14823761

10. Baur A, Harrer T, Peukert M, et al. Alpha-lipoic acid is an effective inhibitor of human immuno-deficiency virus (HIV-1) replication. Klin Wochenschr. 1991 Oct 2:69(15):722-4. [PMID: 1724477]

11. Bhattacharya A. Methylselenocysteine: a promising antiangiogenic agent for overcoming drug delivery barriers in solid malignancies for therapeutic synergy with anticancer drugs. Expert Opin Drug Deliv. 2011 Jun;8(6):749-63. [PMID: 21473705]

12. Shilo S, Pardo M, Aharoni-Simon M, et al. Selenium supplementation increases liver MnSOD expression: molecular mechanism for hepato-protection. J Inorg Biochem. 2008 Jan:102(1):110-8. [PMID: 17804075]

13. Brown MD. Green tea (Camellia sinensis) extract and its possible role in the prevention of cancer. Altern Med Rev. 1999 Oct;4(5):360-70. Review. [PMID: 10559550]

14. Shankar S, Ganapathy S, Srivastava RK. Green tea polyphenols: biology and therapeutic implications in cancer. Front Biosci. 2007 Sep 1;12:4881-99. Review. [PMID: 17569617

15. Ferrer P. Asensi M. Segarra R. et al. Association between pterostilbene and ouercetin inhibits metastatic activity of B16 melanoma. Neoplasia. 2005 Jan:7(1):37-47. [PMID: 15736313] 16. Jurenka JS. Anti-inflammatory properties of curcumin, a major constituent of Curcuma longa: a review of preclinical and clinical research. Altern Med Rev. 2009 Jun; 14(2): 141-53. [PMID: 19594223]

17. Choi H, Chun YS, Shin YJ, et al. Curcumin attenuates cytochrome P450 induction in response to 2,3,7,8-tetrachlorodibenzo-p-dioxin by ROS-dependently degrading AhR and ARNT. Cancer Sci. 2008 Dec;99(12):2518-24. [PMID: 19018768]

18. Shoba G, et al. Influence of piperine on the pharmacokinetics of curcumin in animals and human volunteers. *Planta Med.* 1998 May; 64 (4):353-6. [PMID: 9619120] 19. Dalessandri KM, Firestone GL, Fitch MD, et al. Pilot study: effect of 3,3'-diindolylmethane supplements on urinary hormone metabolites in postmenopausal women with a history of

early-stage breast cancer. Nutr Cancer. 2004;50(2):161-7. [PMID: 15623462]

20. Kim EJ, Shin M, Park H, et al. Oral administration of 3.3'-diindolvlmethane inhibits lung metastasis of 4T1 murine mammary carcinoma cells in BALB/c mice. J Nutr. 2009 Dec:139(12):2373-9. [PMID: 19864400]

21. Bolton JL, Thatcher GR. Potential mechanisms of estrogen quinine carcinogenesis. Chem Res Toxicol. 2008 Jan;21(1):93-101. Review. [PMID: 18052105]

22. Keum YS. Regulation of the Keap1/Nrf2 system by chemopreventive sulforaphane: implications of posttranslational modifications. Ann N Y Acad Sci. 2011 Jul;1229:184-9. Review. [PMID: 21793854]

23. Boddupalli S, Mein JR, Lakkanna S, et al. Induction of phase 2 antioxidant enzymes by broccoli sulforaphane: perspectives in maintaining the antioxidant activity of vitamins A, C, and E. Front Genet. 2012;3:7. Epub 2012 Jan 24. [PMID: 22303412] 24. Sulforaphane glucosinolate. Monograph. Altern Med Rev. 2010 Dec;15(4):352-60. Review. [PMID: 21194251] p. 352.

Formulated To Exclude

Wheat, gluten, yeast, soy, animal and dairy products, fish, shellfish, peanuts, tree nuts, egg, ingredients derived from genetically modified organisms (GMOs), artificial colors, artificial sweeteners, and artificial preservatives.

> *These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

> > Distributed By: MSW Nutrition 3930 Bee Caves RD, STE F Austin, TX 78746 512-356-9144

Directions

Take two capsules daily, or as directed by your healthcare professional.

Consult your healthcare professional prior to use. Individuals taking medication should discuss potential interactions with their healthcare professional. Do not use if tamper seal is damaged.

MITOCHONDRIAC

Clinical Applications

- Methylation helps improve absorption and stability*
- Powerful Antioxidant Support*
- Supports Healthy Cellular Function*
- Produces Changes Associated with Longer Lifespan (including activation of sirtuins)*
- Supports Cardiovascular/Neurological Health*

Mitochondriac provides a complete bioflavonoid complex with resveratrol, quercetin, and pterostilbene. These three antioxidants work together synergistically. Pterostilbene, a new focus of scientific research, is methylated resveratrol, a bio-optimized form that complements resveratrol and quercetin. These compounds are being extensively studied in the areas of cardiovascular health and aging.*

All Mendoza Sports & Wellness Formulas Meet or Exceed cGMP Quality Standards

Discussion

Resveratrol (RES) (3,5,4'-trihydroxystilbene) is a stilbenol derived from stilbene, a natural plant product. RES is found in varying amounts in grapes, various berries, plums, peanuts (and pines). Oral resveratrol is rapidly metabolized via sulfate conjugation by the intestine/liver. ^[1] The methyl capping of all free hydroxyl groups (as in Pterostilbene) results in dramatically higher hepatic metabolic stability, intestinal absorption and membrane transport compared to unmethylated RES.^[2,3] Quercitin, pterostilbene and resveratrol are synergistic antioxidants, with quercitin seemingly aiding in the absorption of resveratrol.^{*[4]}

Stilbenols are polyphenolic compounds that have cell-protecting properties.^[6] For RES, this action has mostly been linked to growth and death regulatory pathways. Research published in 2008 also demonstrated RES's contribution to the maintenance of genome stability. Pterostilbene and quercetin, structurally-related and naturally-occurring, small polyphenols, show longer half-life in vivo than unmethylated resveratrol and have been shown to work synergistically to protect cellular health.*^[6]

As exciting as its role in chemoprotection, is resveratrol's ability to produce changes associated with longevity. These include increased insulin sensitivity, reduced IGF-1, increased AMP-activated protein kinase and peroxisome proliferator-activated receptor-gamma coactivator 1 alpha activity, increased mitochondrial number and improved motor function. RES opposed the effects of the high calorie diet in 144 of 153 significantly altered gene pathways.^[7] RES activates sirtuins including SIR2, a special longevity cellular^[8] and SIRT1 that helps protect nerve cells.^{*[9]}

In vitro, ex vivo and animal experiments have shown that the attributes of RES such as its powerful antioxidant activity, modulation of hepatic apolipoprotein and lipid synthesis, inhibition of platelet aggregation, and inhibition of human platelet and neutrophil production of proatherogenic eicosanoids favor protection against atherosclerosis.*^[10]

Resveratrol's numerous anti-inflammatory properties may explain why it has so many far-reaching health benefits. It inhibits synthesis and release of pro-inflammatory mediators, modifies eicosanoid synthesis, and inhibits activated immune cells. By inhibiting either NF-(kappa)B or the activator protein-1 (AP-1), resveratrol also appears to inhibit inducible nitric oxide synthase (iNOS) and cyclooxygenase-2 (COX-2).^[11] Specific human dosing to support a healthy inflammatory response has not yet been established.^{*[12]}

IITOCHONDRI/



Supplement Facts

Serving Size: 1 Capsule

ourvings for ountainer. oo		
	Amount Per Serving	%Daily Value
Quercetin (as quercetin dihydrate)(from <i>Sophora japonica</i>)(bud)	250 mg	**
<i>trans</i> -Resveratrol (as <i>Polygonum cuspidatum</i> root extract)	75 mg	**
trans-Pterostilbene ^{S1}	62.5 mg	**
** Daily Value not established.		

Other Ingredients: HPMC (capsule), microcrystalline cellulose, stearic acid, magnesium stearate, silica, and medium-chain triglyceride oil.

PTER®PUR® S1. Pteropure® is a registered trademark of ChromaDex, Inc.



Directions

Take one capsule twice daily, or as directed by your healthcare professional.

Consult your healthcare professional prior to use. Individuals taking medication should discuss potential interactions with their healthcare professional. Do not use if tamper seal is damaged.

Storage:

Keep closed in a cool, dry place out of reach of children.

References

1. Wenzel E, Somoza V. Metabolism and bioavailability of trans-resveratrol. Mol Nutr Food Res. 2005 May;49(5):472-81. [PMID:15779070]

2. Walle T, et al. High absorption but very low bioavailability of oral resveratrol in humans. Drug Metab Dispos. 2004 Dec;32(12):1377-82. [PMID:15333514]

3. Wen X, Walle T. Methylated flavonoids have greatly improved intestinal absorption and metabolic stability. Drug Metab Dispos. 2006 Oct;34(10):1786- 92 [PMID: 16868069]

4. Morimitsu Y, Sugihara N, Furuno K. Inhibitory effect of flavonoids on sulfo- and glucurono-conjugation of acetaminophen in rat cultured hepatocytes and liver subcellular preparations. Biol Pharm Bull. 2004 May;27(5):714-7. [PMID: 15133252]

5. Fresco P, Borges F, Diniz C, Marques MP. New insights on the anticancer properties of dietary polyphenols. Med Res Rev. 2006 Nov;26(6):747-66 [PMID: 16710860]

6. Ferrer P, et al. Association between pterostilbene and quercetin inhibits metastatic activity of B16 melanoma. Neoplasia. 2005 Jan;7(1):37-47. [PMID:15736313]

7. Baur JA, et al. Resveratrol improves health and survival of mice on a highcalorie diet. Nature. 2006 Nov; 16;444(7117):337-42. [PMID: 17086191]

8. Stefani M, et al. The effect of resveratrol on a cell model of human aging. Ann N Y Acad Sci. 2007 Oct;1114:407-18. [PMID: 17804521]

9. Tang BL, Chua CESIRT1 and neuronal diseases. Mol Aspects Med. 2008 Jun;29(3):187-200 [PMID: 17397914]

10. Soleas GJ, Diamandis EP, Goldberg DM. Resveratrol: a molecule whose time has come? And gone? Clin Biochem. 1997 Mar;30(2):91-113 [PMID:9127691]

11. de la Lastra CA, Villegas I.Resveratrol as an anti-inflammatory and antiaging agent: mechanisms and clinical implications. Mol Nutr Food Res. 2005 May;49(5):405-30 [PMID:15832402]

12. Hougee S, Faber J, Sanders A, de Jong RB, van den Berg WB, Garssen J, Hoijer MA, Smit HF. Selective COX-2 inhibition by a Pterocarpus marsupium extract characterized by pterostilbene, and its activity in healthy human volunteers. Planta Med. 2005 May;71(5):387-92. [PMID: 15931573] Additional references available upon reques

Formulated To Exclude

Wheat, gluten, yeast, soy, animal and dairy products, fish, shellfish, peanuts, tree nuts, egg, ingredients derived from genetically modified organisms (GMOs), artificial colors, artificial sweeteners, and artificial preservatives.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

> Mendoza Sports & Wellness 3930 Bee Caves RD, STE F West Lake Hills, TX 78746 512-461-9949

Clinical A lications

- Promotes Energy Production and Stamina*
- Supports the Body's Adaptogenic Response*
- Supports the Body's Response to Stress*
- Promotes Adrenal Physiological Functions*

Zen features a comprehensive blend of nutrients and botanical extracts targeted to supporting the body's adaptogenic response to promote optimal energy production, stamina, and the management of everyday stressors. Adrenal glandular tissue, sourced from Argentinian bovine to safeguard purity, rounds out the ingredient profile.*

All MSW Nutrition Formulas Meet or Exceed cGMP Quality Standards

iscussion

Cortisol is one of the main stress-mediating glucocorticoid hormones producedby the adrenal gland. acts lt to maintain blood glucose levels necessary fornormal brain and physical functions. Typically, cortisol levels spike in themorning and gradually decrease throughout the day. When the body is understress, cortisol production increases. With prolonged increased levels of cortisol, the body produces more leading to an glucose increase in risk for depressedimmune function, increased production of cytokines, insulin resistance, weightgain. and other potentially chronic conditions. Eventually, the adrenal productionof cortisol weakens resulting in "adrenal fatique." which causes feelings offiredness, mild depression, and general malaise. Zen is formulated tosupport Individuals who present with cortisol levels that are low throughout theday and are sometimes referred to as being in the "burnout" or "flat-line" stageof adrenal fatigue.*

Vitamin C

The release of adrenocorticotropic hormone (ACTH) from the pituitary gland in tandem with the body's physiological response to stress will deplete the relatively large amount of vitamin C typically stored in the adrenal cortex.^[1,2] This vitamin is essential for the synthesis of epinephrine, the hormone secreted by the adrenal medulla in response to stress. Epinephrine, in turn, plays a role in the synthesis of aldosterone, the hormone that regulates blood pressure, volume, and pH.*

Vitamin B6, Pantothenic Acid

Vitamin B6, acting as a coenzyme, has a role in the conversion of muscle glycogento glucose, which is needed for a proper response to stressors, the synthesis ofserotonin, and the support of the immune function.^[3] Physiologically, vitamin **B6influences** the adrenal glucocorticoid receptor, stimulates the and aids in sodium and potassium balance.*[2]Pantothenic secretion of adrenalcatecholamines, acid is essential to the adrenal glands for the production ofglucocorticoids. It forms pantethine in the body, which then converts tocoenzyme-A-the most active metabolic enzyme in the human body needed toproduce cellular energy.[4] Pantothenic acid plays a critical role in the utilizationof fats and carbohydrates in energy production and in the manufacture of adrenal hormones and red blood cells.*[5]

Asian Ginseng (Panax)

As an important herbal remedy in traditional Chinese medicine, Panax ginseng has been used for thousands of years, primarily for energy production. The main active agents have been identified as ginsenosides, and they are the focus of much published research.^[6] Experimental models show that ginseng and ginsenosides have beneficial effects in supporting the adrenal glands; protecting the gastric mucosa; and supporting healthy body weight, blood hormones, and the gene expression of catecholamine-synthesizing enzymes.^[6-10] Ginsenosides also have immune-supporting and cytokine-modulating activities.^{*[7,11]}

Continued on next page

Distributed By: **MSW Nutritior** 3930 Bee Caves RD, STE F Austin, TX 78746 512-356-9144

Eleuthero

In the 1950s, Russian researchers investigated the properties of eleuthero anddiscovered that it had "adaptogenic" activity. An adaptogen refers to a substancethat supports the body's ability to adapt and promote healthy physiologicalfunctioning, most notably in relation to stress. This balancing effect has beenevaluated in numerous human clinical trials that demonstrate the ability of eleuthero to increase stamina, mental alertness, and the capability of theparticipants to handle stress.^{*[12,19]}

Rhodiola

This adaptogenic herb has been used traditionally in Eastern Europe and Asiafor centuries to increase stamina, maintain a healthy mood, support the nervousand immune systems, and maintain healthy male sexual function.^[13,14] Accordingto Panossian et al, experimental studies performed on isolated organs, tissues,cells, and enzymes demonstrated that rhodiola preparations exhibit adaptogeniceffects that support nerve, brain, and heart health and instill calm, enhancelongevity, and stimulate the nervous system.^[14] Rhodiola may also have apositive effect on brain neurotransmitters, such as dopamine and serotonin,and may influence endogenous opioid levels.^[13] According to a review of theliterature on rhodiola, supplementation supports healthy work performance,quality of sleep, appetite, and energy levels subsequent to intense physical orintellectual strain. Salidrosides and rosavin have been identified as primaryactives. The rhodiola extract in Zen is standardized to provide no less than 1% salidrosides and 3% rosavin.*

Schisandra

Schisandra has been widely studied for its effect on helping the body adapt topsychological and physical stressors. It has been suggested that schisandraaffects the basal levels of nitric oxide and cortisol, which promote enduranceand accuracy of movement, mental performance, and working capacity.*[15,16]

Schisandra is commonly used in combination with other adaptogens. In adouble-blind placebo-controlled trial with rhodiola and eleuthero, improvementin attention as well as speed and accuracy on cognitive tasks was noted.*[17]

Adrenal Gland (from Argentina bovine), Licorice Root

Adrenal gland derived from pure Argentinian bovine is a complementaryingredient traditionally used to replenish adrenal function.^[18] Licorice containstriterpenoid saponins that influence cortisol balance, and it is a staple herb forsupporting adrenal insufficiency.^{*[19]}



Supplement Facts

Serving Size: 1 Capsule Servings Per Container: 60

	Amount Per Serving	%Daily Value
Vitamin C (ascorbic acid)	125 mg	139%
Vitamin B6 (as pyridoxal 5'-phosphate)	15 mg	882%
Pantothenic Acid (as d-calcium pantothenate)	150 mg	3,000%
Adrenal Gland (from bovine)(Argentina)	125 mg	* *
Asian Ginseng Extract (<i>Panax ginseng</i>)(root) (4% ginsenosides)	75 mg	**
Eleuthero Extract (<i>Eleutherococcus</i> <i>senticosus</i>)(root)(0.8% eleutheroside)	50 mg	* *
Licorice (Glycyrrhiza glabra)(roots)	50 mg	**
Schisandra Extract (<i>Schisandra chinensis</i>) (berries)(2% schisandrins)	50 mg	**
Rhodiola Extract (<i>Rhodiola rosea</i>)(root)(3% rosavins and 1% salidroside)	50 mg	* *
** Daily Value not established.		

Other Ingredients: Capsule (hypromellose and water), ascorbyl palmitate, mediumchain triglyceride oil, and silica.



References

1. Murray RK, Granner DK, Mayes PA, et al. Harper's Biochemistry. 25th ed.Stamford, CT: Appleton & Lange; 2000.

2. Head KA, Kelly GS. Nutrients and botanicals for treatment of stress: adrenalfatigue, neurotransmitter imbalance, anxiety, and restless sleep. Altern MedReview. 2009 June;14(2): 114-140. [PMID: 19594222]

3. Vitamin B5 (Pantothenic acid). University of Maryland Medical Center. http://www.umm.edu/health/medical/altmed/supplement/vitamin-b5-pantothenicacid.Updated July 16, 2013. Accessed September 12, 2017.

4. National Institutes of Health. Office of Dietary Supplements. Dietary SupplementFact Sheet: Vitamin B6. https://ods.od.nih.gov/factsheets/VitaminB6-HealthProfessional/ Updated February 11, 2016. Accessed September 11, 2017.

5. Kelly CJ. Invigorating the context and content of nutrition in medical education. Aca Med. 2011 Nov;86(11):1340. [PMID: 22030639]

6. Kiefer D, Pantuso T. Panax ginseng. Am Fam Physician. 2003 Oct15;68(8):1539-42. [PMID: 14596440]

7. Jia L, Zhao Y, Liang XJ. Current evaluation of the millennium phytomedicineginseng (II): Collected chemical entities, modern pharmacology, and clinicalapplications emanated from traditional Chinese medicine. Curr Med Chem.2009;16(22):2924-42. [PMID: 19689273]

8. Kim Y, Choi EH, Doo M, et al. Anti-stress effects of ginseng via down-regulationof tyrosine hydroxylase (TH) and dopamine β-hydroxylase (DBH) geneexpression in immobilizationstressed rats and PC12 cells. Nutr Res Pract. 2010Aug;4(4):270-75. [PMID: 20827341]

9. Tachikawa E, Kudo K, Hasegawa H, et al. In vitro inhibition of adrenalcatecholamine secretion by steroidal metabolites of ginseng saponins. BiochemPharmacol. 2003 Dec 1;66(11):2213-21. [PMID: 14609746]

10. Rai D, Bhatia G, Sen T, et al. Anti-stress effects of Gingko biloba and Panaxginseng: a comparative study. J Pharmacol Sci. 2003 Dec;93(4):458-64. [PMID:14737017]

11. Lee DC, Lau AS. Effects of Panax ginseng on tumor necrosis factor-α-mediatedinflammation: a mini-review. Molecules. 2011 Mar 30;16(4):2802-16. [PMID:21455094] 12. Panossian A, Wikman G. Effects of adaptogens on the central nervous systemand the molecular mechanisms associated with their stress-protective activity.Pharmaceuticals. 2010

Jan 19;3(1):188-224. [PMID: 27713248]

13. Chan SW. Panag Rhodiola rosea and Schisandra chinensis. Int J FoodSci Nutr. 2012 Mar:63 Suppl 1:75-81. [PMID: 22039930]

14. Panossian A, Wikman G, Sarris J. Rosenroot (Rhodiola rosea): traditional use, chemical composition, pharmacology and clinical efficacy. Phytomedicine. 2010Jun;17(7):481-93. [PMID: 20378318]

15. Panossian A, Wikman G. Pharmacology of Schisandra chinensis Bail.: anoverview of Russian research and uses in medicine. J Ethnopharmacol. 2008July;118(2):183-212. [PMID: 18515024]

16. Upton R, ed. Schisandra Berry (Schisandra chinensis): Analytical, Quality Controland Therapeutic Monograph. Santa Cruz, CA: American Herbal Pharmacopoeia;1999:1-25. http:// cms.herbalgram.org/herbclip/pdfs/011007-185.pdf.

17. Aslanyan G, Amroyan E, Gabrielyan E, et al. Double-blind, placebo-controlled, randomised study of single dose effects of ADAPT-232 on cognitive functions. Phytomedicine. 2010 June; 17(7):494-499. [PMID: 20374974]

18. Ronzio R. A rationale for the use of glandular products. Stichting Sci Nutrition.http://tuesdayminute.net/A-Rationale_for_the_use_of_glandular_products.pdf.Revised October 2006. Accessed September 11, 2017.

19. Panossian AG. Adaptogens: Tonic herbs for fatigue and stress. AlternComplement Ther. 2004 July;9(6):327-331. https://www.researchgate.net/publication/244889830_Adaptogens_Tonic_Herbs_for_Fatigue_and_Stress.Accessed September 19, 2017.

Formulated To Exclude

Wheat, gluten, yeast, soy, dairy products, fish, shellfish, peanuts, tree nuts, egg, ingredients derived from genetically modified organisms (GMOs), artificial colors, artificial sweeteners, and artificial preservatives.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.

> Distributed By: **MSW Nutrition** 3930 Bee Caves RD, STE F Austin, TX 78746 512-356-9144

Directions:

Take one capsule twice daily, or as directed by your healthcare professional.

Consult your healthcare professional prior to use. Individuals taking medication should discuss potential interactions with their healthcare professional. Do not use if tamper seal is damaged.

Storage:

Keep closed in a cool, dry place out of reach of children.