

# BLISS



## Clinical Applications

- Supports Biochemical Reactions Requiring Methyl Groups\*
- Supports Neurotransmitter Synthesis and Healthy Mood\*
- Facilitates Conversion of Homocysteine to Glutathione\*
- Supports Liver Health and Function\*
- Promotes Joint Comfort\*

***BLISS** is a sweet, yet slightly tart lemon-flavored powder. **BLISS** is encapsulated SAME. SAME (S-adenosyl-L-methionine) and TMG (trimethylglycine) are naturally occurring substances that act as methyl donors during vital biochemical processes in the body. Methylation is essential to normal cell health and function. It can decline with age or chronic alcohol consumption, and it can be limited in some individuals due to their genetic makeup. SAME donates methyl groups, which are needed for the synthesis of neurotransmitters, proteins, nucleic acids, and phospholipids. It supports glutathione production, liver health, joint comfort, and a healthy mood. TMG is another methyl donor. It is involved in the metabolism of homocysteine and the formation of SAME. TMG ultimately supports cardiovascular and neurological health, as well as normal cell-life regulation.\**

All MSW Nutrition Formulas Meet or Exceed cGMP Quality Standards

## Discussion

S-adenosyl-L-methionine (SAME) is a naturally occurring substance formed in the body from the amino acid methionine and the “energy molecule” adenosine triphosphate (ATP). Formation of SAME is catalyzed by methionine adenosyltransferase and depends on cofactors including vitamin B6, vitamin B12, folate, and magnesium. SAME has been studied as a supportive nutrient in liver health, joint comfort, metabolic reactions, and healthy mood.\*<sup>[1-4]</sup>

**Methylation** SAME is the “universal” methyl donor for biochemical reactions throughout the body.<sup>[5]</sup> This methyl transfer, or “transmethylation,” is critical to reactions involving proteins, phospholipids, DNA, RNA, creatine, hormones, development of cell membranes, degradation of histamine, and formation of norepinephrine and dopamine.<sup>[1,6]</sup> Eighty-five percent of transmethylation takes place in the liver, and healthy SAME levels appear to be essential to liver health and function.\*<sup>[2,7]</sup>

**Antioxidant and Liver Support** SAME is considered to be “critical” for synthesis of glutathione, a principal component of antioxidant and detoxification systems in the body.<sup>[1]</sup> Following donation of a methyl group, SAME is converted to S-adenosyl-homocysteine (SAH). This biochemical reaction promotes the transsulfuration pathway in the liver that generates glutathione. Further metabolism of SAH involves trimethylglycine (TMG), also known as betaine anhydrous. TMG plays an important role in maintaining a healthy SAME:SAH ratio in the liver.\*

During a national symposium, the roles of SAME and TMG in supporting liver health were reviewed with a focus on their participation in the vital processes of transmethylation and transsulfuration, their ultimate contribution to increased glutathione synthesis and its hepatoprotective effects, their promotion of a balanced SAME:SAH ratio, their activation of phosphatidylethanolamine methyltransferase, and the increase in phosphatidylcholine synthesis as a result of their administration.<sup>[6]</sup> Ongoing animal studies suggest that SAME supports liver health<sup>[9,10]</sup> and that exogenous SAME may positively affect cell-life regulation of hepatocytes.<sup>[7]</sup> In certain human cohorts, researchers recommend further research into combining SAME with nutrients such as vitamin B6 to optimize outcomes.\*<sup>[5]</sup>

**Healthy Mood** Supplemental SAME appears to support a healthy mood, possibly due to its active role in methylation and its involvement in the formation of monoamine neurotransmitters.<sup>[3,11-13]</sup> Meta-analysis of earlier studies suggested that SAME showed greater support of a healthy mood when compared to placebo with an effect comparable to that of other treatments.<sup>[14]</sup> A 30-day, double-blind, placebo-controlled, randomized study of 80 women suggested that there was a significant improvement in mood after the women received an oral dose of 1600 mg/d of SAME compared to placebo.<sup>[15]</sup> Another study of 143 subjects who received an oral dose of 1600 mg/d of SAME suggested that SAME yielded positive results that were comparable to other treatments for supporting a healthy mood, but SAME was better tolerated.<sup>[16]</sup> In a small (N=26), four-week, double-blind, randomized protocol comparing oral SAME with other treatments, 62% of the SAME group showed significant improvement in mood. The study revealed a significant correlation between plasma SAME levels and the degree of healthy mood support, regardless of treatment type.\*<sup>[17]</sup>

**TMG** Trimethylglycine is a naturally occurring compound (glycine attached to three methyl groups) that is found in food (estimated intake 0.5-2 g/d) and can be produced in the body from the precursor choline.<sup>[18]</sup> TMG is thought to protect liver cells, support homocysteine metabolism and cardiovascular health, and may also support a healthy mood due to its role in SAME metabolism.<sup>[3,9,18,19]</sup> When TMG donates a single methyl group, it is converted to dimethylglycine (DMG), which is capable of donating two methyl groups. TMG is thought to stimulate activity of the enzyme betaine-homocysteine methyltransferase (BHMT). BHMT, found in abundance in a healthy liver,<sup>[20,21]</sup> is used by TMG to donate a methyl group to homocysteine. Once TMG adds a methyl group to homocysteine to produce methionine, the methionine can then be converted to SAME. A randomized, double-blind, crossover study of healthy volunteers suggested that TMG supplementation (at doses of 3 g and 6 g/d) has a dose-dependent effect on serum TMG levels and a significantly positive effect on maintaining healthy homocysteine levels.<sup>[22]</sup> Together, BLISS & TMG provide an abundant source of methyl groups and ultimately support a wide variety of biochemical reactions in the body.\*

**BLISS** MSW Nutrition’s BLISS contains a minimum of 70% of the SS isomer of SAME, the form the body can use most readily. This relatively high concentration from Gnosis’ Adomix® not only makes BLISS particularly bioavailable but also cost-effective. Each capsule is sealed in a nitrogen-purged blister pack to maximize protection from the environment.\*

**\*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.**

Distributed By: **MSW Nutrition**  
3930 Bee Caves RD, STE F  
Austin, TX 78746  
512-356-9144

**BLISS**



BLISS

# Supplement Facts

Serving Size: 1 Stick Pack (about 2.6 g)  
Servings Per Container: 30

	Amount Per Serving	%Daily Value
Calories	5	
Total Carbohydrate	1 g	<1%†
Calcium (as calcium carbonate, calcium oxide, and calcium chloride anhydrous)	75 mg	6%
Betaine Anhydrous (trimethylglycine)	600 mg	**
S-adenosyl-L-methionine (as s-adenosyl-L-methionine 1,4-butanedisulfonate)	400 mg	**

†Percent Daily Values are based on a 2,000 calorie diet.  
\*\* Daily Value not established.

**Other Ingredients:** Xylitol, citric acid, malic acid, stearic acid, silica, natural lemon flavor, and turmeric extract (for color).

## Directions

Consume the contents of one stick pack daily away from meals, or as directed by your healthcare professional. Preferably pour a small amount of the contents of a stick pack directly into the mouth and allow contents to dissolve. Then repeat process until contents of the entire stick pack have dissolved in the mouth. Alternatively, contents may be added to 2-4 oz of water or preferred liquid; stir and drink within 15 minutes.

Consult your healthcare professional prior to use. Individuals taking medication should discuss potential interactions with their healthcare professional. Use special caution in individuals with bipolar disorder. Do not use if stick pack is damaged.

## References

1. Natural Standard Database. SAME. <http://naturalstandard.com/databases/herbssupplements/same.asp?#undefined>. Accessed October 21, 2012.
2. Friedel HA, Goa KL, Benfield P. S-adenosyl-L-methionine. A review of its pharmacological properties and therapeutic potential in liver dysfunction and affective disorders in relation to its physiological role in cell metabolism. *Drugs*. 1989 Sep;38(3):389-416. [PMID: 2680435]
3. Papakostas GI, Cassiello CF, Iovieno N. Folate and s-adenosylmethionine for major depressive disorder. *Can J Psychiatry*. 2012 Jul;57(7):406-13. [PMID: 22762295]
4. Mayo Clinic. SAME. Updated September 1, 2012. [http://www.mayoclinic.com/health/same/NS\\_patient-same/DSECTION=dosing](http://www.mayoclinic.com/health/same/NS_patient-same/DSECTION=dosing). Accessed October 22, 2012.
5. Halsted CH, Medici V. Vitamin-dependent methionine metabolism and alcoholic liver disease. *Adv Nutr*. 2011 Sep;2(5):421-7. [PMID: 22332083]
6. Lu SC. S-Adenosylmethionine. *Int J Biochem Cell Biol*. 2000 Apr;32(4):391-5. [PMID: 10762064]
7. Mato JM, Lu SC. Role of S-adenosyl-L-methionine in liver health and injury. *Hepatology*. 2007 May;45(5):1306-12. [PMID: 17464973]
8. Purohit V, Abdelmalek MF, Barve S, et al. Role of S-adenosylmethionine, folate, and betaine in the treatment of alcoholic liver disease: summary of a symposium. *Am J Clin Nutr*. 2007 Jul;86(1):14-24. [PMID: 17616758]
9. Kharbada KK. Alcoholic liver disease and methionine metabolism. *Semin Liver Dis*. 2009 May;29(2):155-65. [PMID: 19387915]
10. Kharbada KK, Rogers DD 2nd, Mailliard ME, et al. A comparison of the effects of betaine and S-adenosylmethionine on ethanol-induced changes in methionine metabolism and steatosis in rat hepatocytes. *J Nutr*. 2005 Mar;135(3):519-24. [PMID: 15735087]
11. Baldessarini RJ. Neuropharmacology of S-adenosyl-L-methionine. *Am J Med*. 1987 Nov 20;83(5A):95-103. [PMID: 3318448]
12. Morgan AJ, Jorm AF. Self-help interventions for depressive disorders and depressive symptoms: a systematic review. *Ann Gen Psychiatry*. 2008 Aug 19;7:13. [PMID: 18710579]
13. Miller AL. The methylation, neurotransmitter, and antioxidant connections between folate and depression. *Altern Med Rev*. 2008 Sep;13(3):216-26. [PMID: 18950248]
14. Bressa GM. S-adenosyl-L-methionine (SAME) as antidepressant: meta-analysis of clinical studies. *Acta Neurol Scand Suppl*. 1994;154:7-14. [PMID: 7941964]
15. Salmaggi P, Bressa GM, Nicchia G, et al. Double-blind, placebo-controlled study of S-adenosyl-L-methionine in depressed postmenopausal women. *Psychother Psychosom*. 1993;59(1):34-40. [PMID: 8441793]
16. Delle Chiaie R, Pancheri P, Scapicchio P. Efficacy and tolerability of oral and intramuscular S-adenosyl-L-methionine 1,4-butanedisulfonate (SAME) in the treatment of major depression: comparison with imipramine in 2 multicenter studies. *Am J Clin Nutr*. 2002 Nov;76(5):1172S-6S. [PMID: 12418499]
17. Bell KM, Potkin SG, Carreon D, et al. S-adenosylmethionine blood levels in major depression: changes with drug treatment. *Acta Neurol Scand Suppl*. 1994;154:15-8. [PMID: 7941961]
18. Olthoff MR, Verhoef P. Effects of betaine intake on plasma homocysteine concentrations and consequences for health. *Curr Drug Metab*. 2005 Feb;6(1):15-22. [PMID: 15720203]
19. Yi EY, Kim YJ. Betaine inhibits in vitro and in vivo angiogenesis through suppression of the NF-κB and Akt signaling pathways. *Int J Oncol*. 2012 Nov;41(5):1879-85. doi: 10.3892/ijo.2012.1616. [PMID: 22940742]
20. Wang JA, Dudman NP, Lynch J, et al. Betaine:homocysteine methyltransferase—a new assay for the liver enzyme and its absence from human skin fibroblasts and peripheral blood lymphocytes. *Clin Chim Acta*. 1991 Dec 31;204(1-3):239-49. [PMID: 1819467]
21. Pellanda H, Namour F, Fofou-Caillierez M, et al. A splicing variant leads to complete loss of function of betaine-homocysteine methyltransferase (BHMT) gene in hepatocellular carcinoma. *Int J Biochem Cell Biol*. 2012 Feb;44(2):385-92. [PMID: 22138536]
22. Schwab U, Törrönen A, Meririnne E, et al. Orally administered betaine has an acute and dose-dependent effect on serum betaine and plasma homocysteine concentrations in healthy humans. *J Nutr*. 2006 Jan;136(1):34-8. Erratum in: *J Nutr*. 2007 Apr;137(4):1124. [PMID: 16365055]

## Formulated To Exclude

Wheat, gluten, yeast, soy, animal and dairy products, fish, shellfish, peanuts, tree nuts, egg, ingredients derived from genetically modified organisms (GMOs), artificial colors, artificial sweeteners, and artificial preservatives.

**\*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.**

Distributed By: **MSW Nutrition**  
3930 Bee Caves RD, STE F  
Austin, TX 78746  
512-356-9144

DRS-239  
REV. 09/01/22